

CHARACTERIZATION OF EXTREMAL VALUED FIELDS

SALIH AZGIN, FRANZ-VIKTOR KUHLMANN, AND FLORIAN POP

ABSTRACT. We characterize those valued fields for which the image of the valuation ring under every polynomial in several variables contains an element of maximal value or a zero of the polynomial.

1. INTRODUCTION

The notion of *extremality* for valued fields was introduced by Yuri Ershov in [4] in connection with valued skew fields which are finite-dimensional over their center. It turns out that the original definition given in that paper (and also in talks given by its author) is flawed in the sense that there are no extremal valued fields except algebraically closed valued fields, and Proposition 2 of that paper is false. We fix this flaw by slightly modifying the definition of extremality; see Definition 2.3 below.

The notion of extremality, restricted to certain classes of polynomials, has since become very useful for the characterization of various properties of valued fields, cf. [6].

In valuation theory and particularly the model theory of valued fields, power series fields and, more generally, maximal fields (valued fields without proper immediate extensions) are usually known to have very good properties. For instance, all of them are henselian, and what is more, algebraically complete. So it seemed likely that all of them are also extremal. Our results in this paper will show that this is not the case.

In the present paper, we obtain the following characterization of extremal valued fields with residue characteristic 0:

Theorem 1.1. *Extremal valued fields with residue characteristic 0 are precisely*

- (i) *henselian valued fields whose value group is a \mathbb{Z} -group, and*
- (ii) *henselian valued fields whose value group is divisible and residue field is large.*

More generally, we prove in Section 3:

Theorem 1.2. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ be a valued field. If \mathcal{K} is extremal, then \mathcal{K} is algebraically complete and*

- (i) *Γ is a \mathbb{Z} -group, or*
- (ii) *Γ is divisible and \mathbf{k} is large.*

Date: 30. 6. 2009.

2000 Mathematics Subject Classification: primary: 12J10; secondary: 12E30.

A major part of this research was done while the authors were attending the o-minimality program at the Fields Institute, Jan.-June 2009. The authors would like to thank the Fields Institute for its support and hospitality. The second author was partially supported by a Canadian NSERC grant and by a sabbatical grant of the University of Saskatchewan.

If K is perfect with $\text{char } \mathbf{k} = \text{char } K$, then also the converse holds.

It remains an open question whether the above characterization also holds in the case of mixed characteristic and in the case of non-perfect valued fields.

See the following section for the definitions of *algebraically complete valued field*, *\mathbb{Z} -group* and *large field*.

The authors would like to thank Sergei Starchenko for pointing out the flaw in the definition of extremality and for providing the first example given in Remark 2.4 below.

2. NOTATIONS AND CONVENTIONS

We assume familiarity with the basic concepts of valued fields and their model theoretic properties. We consider valued fields as three-sorted structures

$$\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$$

where K is the underlying field, Γ is the value group, \mathbf{k} is the residue field, $v : K^\times \rightarrow \Gamma$ is the valuation, with valuation ring

$$\mathcal{O}_v := \{a \in K : v(a) \geq 0\}$$

and maximal ideal $\mathfrak{m}_v := \{a \in K : v(a) > 0\}$ of \mathcal{O} . We have the residue class map

$$\begin{aligned} \bar{\cdot} : \mathcal{O}_v &\rightarrow \mathbf{k} \\ a &\mapsto a + \mathfrak{m}_v \end{aligned}$$

and the *residue characteristic of K* is the characteristic of \mathbf{k} . We often refer to K as the valued field, instead of \mathcal{K} . For a subset Δ of an ordered group Γ we use the notation

$$\gamma < \Delta$$

as a shorthand for $\gamma < \delta$ for all $\delta \in \Delta$. We set $v(0) := \infty > \Gamma$, $\Gamma_\infty := \Gamma \cup \{\infty\}$, $-\infty < \Gamma$ and for $\gamma \in \Gamma$ the intervals $(-\infty, \gamma)$ and (γ, ∞) are defined as usual. We use m, n, \dots to denote elements of \mathbb{N} unless specified otherwise.

Definition 2.1. A valued field $(K, \Gamma, \mathbf{k}; v)$ is called algebraically maximal if it does not admit proper immediate algebraic extensions (that is, extensions which preserve value group and residue field). Since henselizations are immediate algebraic extensions, every algebraically maximal field is henselian.

Definition 2.2. A valued field $(K, \Gamma, \mathbf{k}; v)$ is called algebraically complete if it is henselian and for every finite extension $(L, \Delta, \mathbf{l}; w)$ we have

$$(1) \quad [L : K] = (\Delta : \Gamma)[\mathbf{l} : \mathbf{k}].$$

If equation (1) holds, then the extension is called defectless.

Definition 2.3. A valued field \mathcal{K} is extremal if for every multi-variable polynomial $F(X_1, \dots, X_n)$ over K the set

$$\{v(F(a_1, \dots, a_n)) : a_1, \dots, a_n \in \mathcal{O}_v\} \subseteq \Gamma_\infty$$

has a maximal element.

Remark 2.4. The original definition presented in [4] asks for a maximal element of the set

$$\{v(F(a_1, \dots, a_n) : a_1, \dots, a_n \in K\} \subseteq \Gamma_\infty$$

where K and F are as above. This condition is not satisfied for the polynomial

$$F(X, Y) = X^2 + (XY - 1)^2$$

over the Laurent series field $\mathbb{R}((t))$ as

$$v(F(t^n, t^{-n})) = 2n$$

and $F(a, b) \neq 0$ for all $a, b \in \mathbb{R}((t))$. Hence, Proposition 2 of [4] does not hold for the original definition.

Suppose that K is a valued field which is not algebraically closed. Take a polynomial $f(x) = x^n + a_{n-1}x^{n-1} + \dots + a_0 \in K[x]$ with no zeroes in K and let

$$G(X, Y) = X^n + a_{n-1}X^{n-1}Y + \dots + a_0Y^n$$

be its homogenization. Then the polynomial $F(X, Y) = G(X, XY - 1)$ does not satisfy the above condition.

The following is a consequence of Theorem 1.5 of [6]:

Proposition 2.5. *Every extremal field is algebraically maximal and hence henselian.*

We will later use information about the value groups of extremal fields deduced in this paper to prove the stronger assertion that every extremal field is algebraically complete (see Theorem 3.12).

Definition 2.6. *An ordered group is regular if for each n every open interval that contains n elements contains an n -divisible element. A \mathbb{Z} -group is a regular group which is discrete, i.e., has a smallest positive element.*

Note that \mathbb{Z} -groups are exactly the ordered groups which are elementarily equivalent to \mathbb{Z} . If Γ is dense and regular then for each n and $\gamma \in \Gamma$ there is an increasing sequence $\{n\gamma_\rho\}_{\rho < \lambda}$ which is cofinal in $(-\infty, \gamma)$. We state two useful facts on regular groups, from [9] and [1] respectively.

Theorem 2.7. *A regular group is either a \mathbb{Z} -group or it is dense.*

Theorem 2.8. *An ordered group Γ is regular if and only if Γ/Δ is divisible for every nonzero convex subgroup Δ of Γ .*

Definition 2.9. *Take $\mathcal{A} \subseteq \mathcal{B}$ to be an extension of two structures of a first order language. Then \mathcal{A} is existentially closed in \mathcal{B} if every existential sentence with parameters from \mathcal{A} that holds in \mathcal{B} also holds in \mathcal{A} .*

Definition 2.10. *A field \mathbf{k} is large if it is existentially closed in $\mathbf{k}((t))$.*

Algebraically closed fields, real closed fields, pseudo-algebraically closed fields, fields equipped with a henselian valuation are all large fields. Finite fields are not large.

For the following result, see Theorem 17 of [5].

Theorem 2.11. *Suppose that \mathbf{k} is a large and perfect subfield of a field F . If \mathbf{k} is the residue field of a valuation on F , then \mathbf{k} is existentially closed in F (as a field).*

Finally, we will need the following two well known technical lemmas.

Lemma 2.12. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ be a valued field such that $v = w \circ \bar{w}$. Then \mathcal{K} is henselian if and only if w is henselian on K and \bar{w} is henselian on the residue field Kw of K under w . The same holds for “algebraically complete” in the place of “henselian”.*

By use of Hensel’s Lemma, one proves:

Lemma 2.13. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ be a perfect and henselian valued field. Then every embedding of a subfield of \mathbf{k} such that v is trivial on the image can be extended to an embedding of \mathbf{k} such that v is trivial on the image.*

3. NECESSARY AND SUFFICIENT CONDITIONS FOR EXTREMALITY

The first non-algebraically closed examples of extremal fields were provided by Proposition 2 in [4]:

Theorem 3.1. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ be an algebraically complete valued field with $\Gamma \simeq \mathbb{Z}$. Then \mathcal{K} is extremal.*

By Remark 2.4, the this theorem would be false with the definition of extremality as stated in [4]. However, it is easy to check that the proof of Theorem 3.1 given in [4] is valid with the revised definition of extremality.

Note that extremality is a first order condition for valued fields. Hence we can apply the following Ax-Kochen-Ershov Principle for tame valued fields proved in [7], to show extremality for larger classes of valued fields:

Theorem 3.2. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ and $\mathcal{L} = (L, \Delta, \mathbf{l}; w)$ be two perfect algebraically complete valued fields with $\text{char } \mathbf{k} = \text{char } K$. If $\mathbf{k} \equiv \mathbf{l}$ and $\Gamma \equiv \Delta$, then $\mathcal{K} \equiv \mathcal{L}$. If in addition \mathcal{L} is an extension of \mathcal{K} , and if $\mathbf{k} \prec \mathbf{l}$ and $\Gamma \prec \Delta$, then $\mathcal{K} \prec \mathcal{L}$. The same holds with “ \prec_{\exists} ” in the place of “ \prec ” as soon as \mathcal{K} is perfect and algebraically complete with $\text{char } \mathbf{k} = \text{char } K$.*

Here, “ \equiv ” denotes “elementarily equivalent”, “ \prec ” denotes “elementary extension” and “ \prec_{\exists} ” denotes “existentially closed in”.

As a corollary to Theorems 3.1 and 3.2, we obtain:

Theorem 3.3. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ be a perfect algebraically complete valued field with Γ a \mathbb{Z} -group, and $\text{char } \mathbf{k} = \text{char } K$. Then \mathcal{K} is extremal.*

If the residue field has characteristic 0, then “perfect algebraically complete” is equivalent to “henselian”; this is a consequence of the Lemma of Ostrowski.

Theorem 3.4. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ be a perfect algebraically complete valued field with divisible value group Γ and large residue field \mathbf{k} with $\text{char } \mathbf{k} = \text{char } K$. Then \mathcal{K} is extremal.*

Proof. In view of Theorem 3.2, we only have to prove our theorem in the case where Γ is the ordered additive group of the real numbers. Take any polynomial $F \in K[X_1, \dots, X_n]$. Take $\mathcal{K}^* = (K^*, \Gamma^*, \mathbf{k}^*; v^*)$ to be a $|K|^+$ -saturated elementary extension of \mathcal{K} . Then Γ^* is an elementary extension of Γ and \mathbf{k}^* is an elementary extension of \mathbf{k} . Now we distinguish two cases.

Case 1: $\{v(F(a_1, \dots, a_n)) : a_1, \dots, a_n \in \mathcal{O}_v\} \setminus \{\infty\}$ is cofinal in Γ . We wish to show that F has a zero in \mathcal{O}_v^n .

By our choice of \mathcal{K}^* there are $a_1^*, \dots, a_n^* \in \mathcal{O}_{v^*}$ such that $v^*(F(a_1^*, \dots, a_n^*)) > \Gamma$. Hence there exists a valuation w on \mathcal{K}^* that is coarser than v^* , satisfies $wF(a_1^*, \dots, a_n^*) > 0$, but is trivial on K . So we can consider K as a subfield of the residue field K^*w of K^* under w . We write $v^* = w \circ \bar{w}$ where \bar{w} is the valuation induced by v^* on K^*w . Note that $a_1^*, \dots, a_n^* \in \mathcal{O}_w$. We obtain that $0 = F(a_1^*, \dots, a_n^*)w = F(a_1^*w, \dots, a_n^*w)$ and that $a_1^*w, \dots, a_n^*w \in \mathcal{O}_{\bar{w}}$. Since Γ is divisible, $\Gamma \prec_{\exists} \Gamma'$. Further, \mathbf{k}^* is equal to the residue field of K^*w under \bar{w} , and $\mathbf{k} \prec_{\exists} \mathbf{k}^*$. Hence by Theorem 3.2, $\mathcal{K} \prec_{\exists} (K^*w, \Gamma', \mathbf{k}^*; \bar{w})$. Since F has a zero in the valuation ring of the latter field, this implies that there are $a_1, \dots, a_n \in \mathcal{O}_v$ such that $F(a_1, \dots, a_n) = 0$.

Case 2: $\{v(F(a_1, \dots, a_n)) : a_1, \dots, a_n \in \mathcal{O}_v\}$ is not cofinal in $\Gamma = \mathbb{R}$. Then there is some real number r which is the supremum of this set. We wish to show that it is a member of the set.

By our choice of \mathcal{K}^* there are $a_1^*, \dots, a_n^* \in \mathcal{O}_{v^*}$ such that $\delta^* := v^*(F(a_1^*, \dots, a_n^*)) > v(F(a_1, \dots, a_n))$ for all $a_1, \dots, a_n \in \mathcal{O}_v$. On the other hand, $\delta^* \leq r$ in Γ^* . So $0 \leq r - \delta^* < s$ for all $s \in \mathbb{R} = \Gamma$. We take Γ_0 to be the largest convex subgroup of Γ^* such that $\Gamma_0 \cap \Gamma = \{0\}$. Then we take w to be the coarsening of v^* with respect to Γ_0 , that is, \mathcal{O}_w is generated over \mathcal{O}_v by all elements whose values under v lie in Γ_0 , and the value group of w on K^* is Γ^*/Γ_0 . Again, we write $v^* = w \circ \bar{w}$ where \bar{w} is the valuation induced by v^* on the residue field K^*w . We observe that $v = w$ on K (after identification of equivalent valuations). Further, $w(F(a_1^*, \dots, a_n^*)) = r$.

Now we have that $(K^*, \Gamma^*/\Gamma_0, K^*w; w)$ is an extension of \mathcal{K} . We have canonical embeddings of \mathbf{k} in K^*w and of Γ in Γ^*/Γ_0 . Since Γ is divisible, $\Gamma \prec_{\exists} \Gamma^*/\Gamma_0$. The residue field K^*w has a valuation \bar{w} with residue field \mathbf{k}^* . Being an elementary extension of K , also K^* is perfect. It follows that its residue fields K^*w and \mathbf{k}^* are perfect. Since every algebraically complete valued field is henselian, Lemma 2.12 shows that K^*w is henselian under \bar{w} . By Lemma 2.13, the canonical embedding of \mathbf{k} in K^*w can be extended to an embedding of \mathbf{k}^* in K^*w . Via this embedding we may assume that \mathbf{k}^* is a subfield of K^*w . Thus by Theorem 2.11, $\mathbf{k}^* \prec_{\exists} K^*w$ and consequently, $\mathbf{k} \prec_{\exists} K^*w$. It now follows from Theorem 3.2 that $\mathcal{K} \prec_{\exists} (K^*, \Gamma^*/\Gamma_0, K^*w; w)$. Hence there are $a_1, \dots, a_n \in \mathcal{O}_v$ such that $vF(a_1, \dots, a_n) = r$. \square

Proposition 3.5. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ be a valued field where Γ is dense, regular and not divisible. Then \mathcal{K} is not extremal.*

Proof. Let $\gamma \in \Gamma$ be positive and not divisible by $n > 1$. Consider the polynomial

$$F(x, y) = x^{4n} + \epsilon(xy - \epsilon^2)^n + \epsilon^2 y^{4n}$$

over K where $v(\epsilon) = \gamma$. Note that for all $a, b \in K^\times$,

$$v(a^{4n}), \quad v(\epsilon(ab - \epsilon^2)^n), \quad v(\epsilon^2 b^{4n})$$

are distinct elements of Γ_∞ since γ is not n -divisible. Hence the valuation of $F(a, b)$ is equal to the minimum of the three distinct elements above.

We claim that for all $a, b \in \mathcal{O}_v$, $v(F(a, b)) < 4n\gamma + \gamma$. Assume otherwise. Then $v(a^{4n}) \geq 4n\gamma + \gamma$ and since equality is not possible,

$$4nv(a) > 4n\gamma + \gamma.$$

Also, $v(\epsilon(ab - \epsilon^2)^n) \geq 4n\gamma + \gamma$ and so $v(a) + v(b) = 2\gamma$. Consequently,

$$4nv(b) = 8n\gamma - 4nv(a)$$

which gives $4nv(b) < 4n\gamma - \gamma$. Then

$$v(F(a, b)) = v(\epsilon^2 b^{4n}) = 2\gamma + 4nv(b) < 4n\gamma + \gamma,$$

contradiction.

Since Γ is dense and regular we can take a sequence $\{4n\delta_\rho\}_{\rho < \lambda}$ in Γ which is cofinal in the interval $(4n\gamma, 4n\gamma + \gamma)$. For each $\rho < \lambda$, pick $a_\rho \in \mathcal{O}_v$ with $v(a_\rho) = \delta_\rho < 2\gamma$ and let $b_\rho := \epsilon^2 a_\rho^{-1} \in \mathcal{O}_v$. Then for all ρ

$$v(a_\rho^{4n}) = 4n\delta_\rho < 4n\gamma + \gamma < v(\epsilon^2 b_\rho^{4n}) = 2\gamma + 4n(2\gamma - \delta_\rho)$$

and hence $v(F(a_\rho, b_\rho)) = 4n\delta_\rho$. Together with the previous claim, this shows that $\{v(F(a, b)) : a, b \in \mathcal{O}_v\}$ has no maximal element and so \mathcal{K} is not extremal. \square

Proposition 3.6. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}, v)$ be a valued field where Γ has a nonzero convex subgroup Δ such that Γ/Δ is not divisible. Then \mathcal{K} is not extremal.*

Proof. Let $\gamma \in \Gamma$ be such that the coset $\gamma + \Delta$ is not divisible by n in Γ/Δ . Consider the polynomial (as introduced in the previous theorem)

$$F(x, y) = x^{4n} + \epsilon(xy - \epsilon^2)^n + \epsilon^2 y^{4n}$$

over K where $v(\epsilon) = \gamma$.

The arguments in the proof of Proposition 3.5 can be applied to the ordered group Γ/Δ to conclude that for all $a, b \in \mathcal{O}_v$,

$$(2) \quad v(F(a, b)) < 4n\gamma + \gamma + \Delta.$$

We claim that $\{v(F(a, b)) : a, b \in \mathcal{O}_v\}$ has no maximal element. Take $a, b \in \mathcal{O}_v$ and set $\theta = v(F(a, b))$. We aim to find $a', b' \in \mathcal{O}_v$ with $v(F(a', b')) > \theta$. Take any positive $\delta \in \Delta$. From (2) it follows that

$$(3) \quad \theta + 4n\delta < 4n\gamma + \gamma - 4n\delta.$$

Note that $v(F(\epsilon, \epsilon)) = 4n\gamma$, so we assume that $4n\gamma < \theta$. As $v(\epsilon) = \gamma$ is not divisible by n , θ is equal to the minimum of $4nv(a)$, $\gamma + nv(ab - \epsilon^2)$ and $2\gamma + 4nv(b)$. Therefore,

$$4n\gamma < \theta \leq v(\epsilon(ab - \epsilon^2)^n)$$

and hence

$$v(b) = 2\gamma - v(a).$$

Assume that $\theta = 4nv(a)$. Together with $4nv(a) < 4n\gamma + \gamma$, (3) shows that

$$(4) \quad 4nv(a) + 4n\delta < \gamma + 4n\gamma - 4n\delta < 2\gamma + 8n\gamma - 4nv(a) - 4n\delta.$$

Now take $a' \in \mathcal{O}_v$ with $v(a') = v(a) + \delta$ and let $b' = \epsilon^2/a' \in \mathcal{O}_v$. Then $v(b') = 2\gamma - v(a) - \delta$ and (4) implies that $4nv(a') < 2\gamma + 4nv(b')$. Therefore $v(F(a', b')) = 4nv(a') = 4nv(a) + 4n\delta > \theta$ as required.

Next assume that $\theta = \gamma + nv(ab - \epsilon^2) \geq 4n\gamma$. Then $\theta < 4nv(a)$ and $\theta < 2\gamma + 4nv(b)$. With $b' = \epsilon^2/a \in \mathcal{O}_v$, we have $v(b') = v(b)$ and therefore $v(F(a, b')) \geq \min\{4nv(a), 2\gamma + 4nv(b')\} > \theta$.

It remains to consider the case $\theta = 2\gamma + 4nv(b)$. Together with $2\gamma + 4nv(b) = \theta < 4n\gamma + \gamma$, (3) shows that

$$\begin{aligned} 2\gamma + 4nv(b) + 4n\delta &< 4n\gamma + \gamma - 4n\delta + (4n\gamma + \gamma - 2\gamma - 4nv(b)) - 4n\delta \\ &= 8n\gamma - 4nv(b) - 4n\delta. \end{aligned}$$

Now take $b' \in \mathcal{O}_v$ with $v(b') = v(b) + \delta$ and let $a' = \epsilon^2/b' \in \mathcal{O}_v$. Then $v(a') = 2\gamma - v(b) - \delta$ and the above inequality implies that $2\gamma + 4nv(b') < 4nv(a')$. Therefore $v(F(a', b')) = 2\gamma + 4nv(b) + 4n\delta > \theta$ as required. This completes the proof of our claim and we conclude that \mathcal{K} is not extremal. \square

Theorem 3.7. *Suppose that Γ is not divisible and not a \mathbb{Z} -group. Then $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ is not extremal.*

Proof. By Theorem 2.7, if Γ is regular then Γ is dense and hence by Proposition 3.5 \mathcal{K} is not extremal. If Γ is not regular then by Theorem 2.8, Γ has a nonzero convex subgroup Δ such that Γ/Δ is not divisible and by Proposition 3.6 \mathcal{K} is not extremal. \square

Proposition 3.8. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ be a valued field with non-algebraically closed residue field \mathbf{k} and divisible value group Γ . Suppose that $f(Y, Z)$ is a polynomial over \mathcal{O}_v such that*

- (i) *there are $\alpha_1, \dots, \alpha_m \in \mathbf{k}$ such that the equation*

$$\bar{f}(\alpha, z) = 0$$

over \mathbf{k} has no solution in \mathbf{k} whenever $\alpha \neq \alpha_1, \dots, \alpha_m$;

- (ii) *for all $\epsilon \in \mathfrak{m}_v$, $f(\epsilon, Z) = 0$ has a solution in \mathcal{O}_v .*

Then \mathcal{K} is not extremal.

Proof. Let $\beta_0, \dots, \beta_{n-1} \in \mathbf{k}$ be such that $x^n + \dots + \beta_1 x + \beta_0 = 0$ has no solutions in \mathbf{k} . Consider the polynomial

$$F(X, Y) = X^n + \dots + b_1 XY^{n-1} + b_0 Y^n$$

over \mathcal{O}_v where $\bar{b}_i = \beta_i$ and $b_i = 0$ if $\beta_i = 0$, for $i = 0, \dots, n-1$. Note that for all positive $\gamma \in \Gamma$,

$$v(F(a, b)) \geq \gamma \implies v(a), v(b) \geq \gamma/n.$$

Let $G(X, Y, Z) := F(F(X, Y), Z)$. For positive $\gamma \in \Gamma$, if $v(G(a, b, c)) \geq \gamma$ then $v(a), v(b) \geq \gamma/n^2$ and $v(c) \geq \gamma/n$.

Let $a_1, \dots, a_m \in \mathcal{O}_v$ be such that $\bar{a}_j = \alpha_j$ and $a_j = 0$ if $\alpha_j = 0$ (where α_j is as in (i) above) for $j = 1, \dots, m$. Consider the polynomial

$$H(X, Y, Z) := G(X, f(Y, Z), XY(Y - a_1) \cdots (Y - a_m) - \epsilon)$$

over K where $v(\epsilon) = \gamma > 0$. We claim that

$$v(H(a, b, c)) < n^2\gamma$$

for all $a, b, c \in \mathcal{O}_v$. Otherwise, we would have

$$v(a) \geq \gamma \text{ and } v(ab(b - a_1) \cdots (b - a_m) - \epsilon) \geq n\gamma.$$

Therefore $v(ab(b - a_1) \cdots (b - a_m)) = v(\epsilon) = \gamma$, which in turn gives

$$v(a) = \gamma, \quad v(b) = 0 \text{ and } v(b - a_i) = 0 \quad \text{for } i = 1, \dots, m.$$

So $\bar{b} \neq \alpha_1, \dots, \alpha_m$ and hence for all $z \in \mathcal{O}_v$, $\bar{f}(\bar{b}, \bar{z}) \neq 0$. Then $v(H(a, b, c)) = v(f(b, c)) = 0$, contradiction, and we establish the claim.

Let $\{\gamma_\rho\}_{\rho < \lambda}$ be a decreasing sequence such that $\gamma_\rho \rightarrow 0$ and pick $b_\rho \in \mathfrak{m}_v$ with $v(b_\rho) = \gamma_\rho$ for all $\rho < \lambda$. Let a_ρ be such that $a_\rho b(b_\rho - a_1) \cdots (b_\rho - a_m) = \epsilon$. Note

that $v(a_\rho) \rightarrow \gamma$, and in particular we may assume that $a_\rho \in \mathcal{O}_v$ for all $\rho < \lambda$. Pick $c_\rho \in \mathcal{O}_v$ such that $f(b_\rho, c_\rho) = 0$ for each $\rho < \lambda$. Then

$$H(a_\rho, b_\rho, c_\rho) = n^2 v(a_\rho)$$

and hence $\{v(H(a_\rho, b_\rho, c_\rho))\}$ is cofinal in $(0, n^2\gamma)$. We conclude that \mathcal{K} is not extremal. \square

Theorem 3.9. *Suppose that $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ has a divisible value group. If \mathbf{k} is not large then \mathcal{K} is not extremal.*

Proof. Since \mathbf{k} is not large, there is a curve C defined over \mathbf{k} which has a smooth \mathbf{k} -rational point, but has only finitely many \mathbf{k} -rational points. But then by the theory of algebraic curves, the curve C is birational to a curve in the affine \mathbf{k} -plane $C_h = V(h) \subset \mathbf{A}_{\mathbf{k}}^2$, where $h = h(Y, Z) \in \mathbf{k}[Y, Z]$ is a polynomial in the variables Y, Z satisfying the following:

- (i)' $C_h(\mathbf{k}) = \{(\alpha, \beta) \in \mathbf{k}^2 \mid h(\alpha, \beta) = 0\}$ is finite.
- (ii)' $h(0, 0) = 0$ and $\partial h / \partial Z(0, 0) \neq 0$, hence in particular, $(0, 0)$ is a smooth \mathbf{k} -rational point of C_h . (Actually these conditions imply that $h(Y, Z)$ has the form $h(Y, Z) = \gamma_{10}Y + \gamma_{01}Z + (\text{non-linear terms})$ with $\gamma_{01} \neq 0$.)

Setting $h(Y, Z) = \sum_{i,j} \gamma_{ij} Y^i Z^j$, let $f(Y, Z) = \sum_{i,j} c_{ij} Y^i Z^j \in \mathcal{O}[Y, Z]$ be a preimage of $h(Y, Z)$ such that $c_{ij} = 0$ if $\gamma_{ij} = 0$, hence $c_{ij} \in \mathcal{O}^\times$ if $\gamma_{ij} \neq 0$. Finally, let $\{\alpha_1 = 0, \dots, \alpha_m\}$ be the set of all the Y -coordinates of the points $(\alpha, \beta) \in C_h(\mathbf{k})$. We claim that $f(Y, Z) \in \mathcal{O}[Y, Z]$ satisfies the conditions (i), (ii) from Proposition 3.8. Indeed, condition (i) is obviously satisfied, because $\bar{f}(Y, Z) = h(Y, Z)$, and $h(Y, Z)$ satisfies condition (i)'. For condition (ii), proceed as follows: For $\epsilon \in \mathfrak{m}_v$ set $f_\epsilon(Z) := f(\epsilon, Z) \in \mathcal{O}[Z]$. Then $\bar{f}_\epsilon(Z) = h(0, Z)$, hence $Z = 0$ is a simple zero of $\bar{f}_\epsilon(Z) = h(0, Z)$, by condition (ii)'. Therefore, by Hensel's Lemma there exists a unique $\eta \in \mathfrak{m}_v$ such that $f_\epsilon(\eta) = 0$, hence equivalently, $f(\epsilon, \eta) = f_\epsilon(\eta) = 0$.

By Proposition 3.8 we conclude that \mathcal{K} is not extremal. \square

The results proved so far enable us to give a version of Proposition 1 of [4] for the modified notion of “extremal field”.

Definition 3.10. *We say that a basis b_1, \dots, b_n of a valued field extension $(K, \Gamma, \mathbf{k}; v) \subset (L, \Delta, l; w)$ is a valuation basis if for all choices of $c_1, \dots, c_n \in K$,*

$$v \sum_{i=1}^n c_i b_i = \min_i v c_i b_i .$$

Note that every finite defectless extension admits a valuation basis.

Lemma 3.11. *Every finite defectless extension of an extremal field is again an extremal field.*

Proof. Take an extremal field $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ and a finite defectless extension $(L, \Delta, l; w)$ of degree m ; we wish to show that the latter is an extremal field. From Theorem 3.7 we know that Γ is divisible or a \mathbb{Z} -group. Therefore, all cosets of Γ in Δ admit representatives that are either 0 or lie between 0 and the smallest positive element of Γ . Consequently, we can choose a valuation basis b_1, \dots, b_m of the

extension such that the values vb_i have the same property. Write $\overline{Y} = (Y_1, \dots, Y_m)$ and take

$$h(\overline{Y}) = N_{L(\overline{Y})|K(\overline{Y})} \left(\sum_{i=1}^m b_i Y_i \right)$$

be the norm form with respect to the basis b_1, \dots, b_m of the extension $L(\overline{Y})|K(\overline{Y})$.

Take a polynomial $F(X_1, \dots, X_n)$ over L ; we wish to show that the set

$$\{v(F(a_1, \dots, a_n)) : a_1, \dots, a_n \in \mathcal{O}_v\}$$

has a maximal element. Denote by $G(\overline{Z})$ the polynomial obtained from F by substituting $\sum_{i=1}^m b_i Z_{ij}$ for X_i , $1 \leq i \leq n$. The polynomial $G(\overline{Z})$ can be written as $G(\overline{Z}) = \sum_{i=1}^m b_i G_i(\overline{Z})$ with $G_i(\overline{Z}) \in K[\overline{Z}]$ for every i . Now let

$$H(\underline{Z}) = h(G_1(\underline{Z}), \dots, G_m(\underline{Z})) = N_{L(\underline{Y})|K(\underline{Y})}(G(\underline{Z})) \in K[\underline{Z}].$$

Since \mathcal{K} is extremal, there exist elements $c_{ij} \in \mathcal{O}_v$, $1 \leq i \leq m$, $1 \leq j \leq n$, such that

$$vH(\overline{c}) = \max\{vH(\overline{c}') \mid c'_{ij} \in \mathcal{O}_v\}.$$

For $1 \leq j \leq n$, we set $d_j = \sum_{i=1}^m b_i c_{ij}$. We wish to show that

$$wF(\overline{d}) = \max\{wF(\overline{d}') \mid d'_j \in \mathcal{O}_w\}.$$

Note that $vN_{L|K}(a) = mva$ for every $a \in L$ since \mathcal{K} is henselian by Proposition 2.5. Take $d'_j \in \mathcal{O}_w$ and write $d'_j = \sum_{i=1}^m b_i c'_{ij}$ with $c'_{ij} \in K$. Since the b_i form a valuation basis,

$$0 \leq vd'_j = v \sum_{i=1}^m b_i c'_{ij} = \min_i vb_i c'_{ij}.$$

Hence for $1 \leq i \leq m$ and $1 \leq j \leq n$, $vb_i + vc'_{ij} \geq 0$. By our assumptions on the values vb_i , this implies that $c'_{ij} \in \mathcal{O}_v$. Now we compute

$$\begin{aligned} wF(\overline{d}') &= \frac{1}{m} vN_{L|K}(F(\overline{d}')) = \frac{1}{m} vH(\overline{c}') \\ &\leq \frac{1}{m} vH(\overline{c}) = \frac{1}{m} vN_{L|K}(F(\overline{d})) = wF(\overline{d}). \end{aligned}$$

□

This lemma yields:

Theorem 3.12. a) *Every extremal valued field is algebraically complete.*

b) *Every finite extension of an extremal field is again an extremal field.*

Proof. a): By Corollary 2.10 of [6], a valued field is algebraically complete if each of its finite defectless extensions is algebraically maximal. By Proposition 2.5, every extremal field is algebraically maximal. Hence part a) of our theorem follows from the preceding lemma.

b): By part a), every finite extension of an extremal field is defectless. Hence the assertion of part b) follows from the preceding lemma. □

Theorem 1.2 now follows from Theorems 3.3, 3.4, 3.7, 3.9 and 3.12.

4. SOME FURTHER RESULTS

An Ax-Kochen-Ershov Principle as in Theorem 3.2 also holds for formally \wp -adic fields (see [8]) and, more generally, for finitely ramified fields (see [3], [10]). A formally \wp -adic field is \wp -adically closed if and only if it is henselian and its value group is a \mathbb{Z} -group. Formally \wp -adic and finitely ramified fields are algebraically complete as soon as they are henselian. Hence, we obtain from Theorems 3.1 and 1.1 via the Ax-Kochen-Ershov Principle:

Theorem 4.1. *A formally \wp -adic field is extremal if and only if it is \wp -adically closed. A finitely ramified field is extremal if and only if it is henselian and its value group is a \mathbb{Z} -group.*

If $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ is a valued field such that $v = w \circ \bar{w}$, then by Lemma 2.12, v is henselian if and only if w and \bar{w} are. The same holds for “algebraically complete” in the place of “henselian”. The corresponding assertion for “extremal” is not entirely known. We leave the easy proof of the following result to the reader:

Lemma 4.2. *In the above situation, if \mathcal{K} is extremal, then K is also extremal with respect to the coarsening v_1 of v .*

From our characterization of extremal fields we obtain:

Proposition 4.3. *Let $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ be a perfect algebraically complete valued field such that $\text{char } \mathbf{k} = \text{char } K$ and $v = v_1 \circ v_2$. Then $\mathcal{K} = (K, \Gamma, \mathbf{k}; v)$ is extremal if and only if K is extremal with respect to v_1 and the residue field of K under v_1 is extremal with respect to v_2 .*

REFERENCES

- [1] Paul F. Conrad. Regularly ordered groups. *Proc. Amer. Math. Soc.*, 13:726–731, 1962.
- [2] Françoise Delon. Quelques propriétés des corps valués en théories des modèles. Thèse Paris VII (1981).
- [3] Yu. L. Ershov. On the elementary theory of maximal valued fields III. *Algebra Logika* **6**:3 (1967), 31–38.
- [4] Yu. L. Ershov. Extremal valued fields. *Algebra Logika*, 43(5): 582–588, 631, 2004.
- [5] F.-V. Kuhlmann. On places of algebraic function fields in arbitrary characteristic. *Advances in Math.* **188** (2004), 399–424.
- [6] F.-V. Kuhlmann. A classification of Artin-Schreier defect extensions and a characterization of defectless fields. to appear
- [7] F.-V. Kuhlmann. The model theory of tame valued fields. In preparation. Preliminary version published in: Structures Algébriques Ordonnées, Séminaire 2007–2008, Paris VII (2009)
- [8] Alexander Prestel and Peter Roquette. Formally p -adic fields. *Lecture Notes Math.* **1050**, Berlin–Heidelberg–New York–Tokyo (1984).
- [9] Abraham Robinson and Elias Zakon. Elementary properties of ordered abelian groups. *Trans. Amer. Math. Soc.*, 96:222–236, 1960.
- [10] M. Ziegler. Die elementare Theorie der henselschen Körper. Inaugural Dissertation, Köln (1972).

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, 1280 MAIN STREET
WEST, HAMILTON, L8S 4K1 ONTARIO, CANADA
E-mail address: `sazgin@math.mcmaster.ca`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SASKATCHEWAN, 106 WIGGINS
ROAD, SASKATOON, SASKATCHEWAN, CANADA S7N 5E6
E-mail address: `fvk@math.usask.ca`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, 209 SOUTH 33RD STREET,
PHILADELPHIA, PA 19104-6395, USA
E-mail address: `pop@math.upenn.edu`